



## Intestinal Parasite Infection and Anemia among Pregnant Women in Ethiopia A Systematic Review and Meta-Analysis

<sup>1</sup>Eleni Tesfaye Tegegne, <sup>2</sup>Mekibib Kassa Tessema, <sup>3</sup>Kaleab Tesfaye Tegegne\*, <sup>3</sup>Abiyu Ayalew Assefa

<sup>1</sup>College of Medicine and Health Science, School of Nursing, University of Gondar, Gondar, Ethiopia

<sup>2</sup>Leishmania Research and Treatment Center, University of Gondar, Gondar, Ethiopia

<sup>3</sup>Department of Public Health, Hawassa College of Health Science, Hawassa, Ethiopia

Submission Date: 01 Aug. 2022 | Published Date: 31 Aug. 2022

\*Corresponding author: Kaleab Tesfaye Tegegne

Department of Public Health, Hawassa College of Health Science, Hawassa, Ethiopia

### Abstract

**Objective:** The aim of this systemic review and meta-analysis was to examine the relationship between intestinal parasitic infection and anemia among pregnant women in Ethiopia

We include six studies in different regions of Ethiopia. We have done this study focusing on intestinal parasitic infection

**Materials and Methods:** The databases searched were PUBMED and Advanced Google Scholar. on reference manager software reporting intestinal parasitic infection and anemia among pregnant women . Three researchers were carried out the data extraction and assessed independently the articles for inclusion in the review using risk-of-bias tool guided by PRISMA checklist. The combined adjusted Odds ratios (OR)) and 95% confidence intervals were calculated using random effect model

**Results:** six observational studies involving 2838 participants, 557 pregnant women who have anemia were included. The combined effect size (OR) for anemia comparing pregnant women who have intestinal parasitic infection versus pregnant women who did not have intestinal parasitic infection was 3.74 (ORMH = 3.74, 95% CI 2.58-5.43) Heterogeneity:  $\tau^2 = 0.13$ ;  $\chi^2 = 13.11$ ,  $df = 5$  ( $P = 0.02$ );  $I^2 = 62\%$  Test for overall effect:  $Z = 6.94$  ( $P < 0.00001$ )

No publication bias was observed (Egger's test:  $p = 0.074$ , Begg's test:  $p = 0.091$  ). 23.99% (681) pregnant women have intestinal parasitic infection during current pregnancy

In all studies, the proportion of anemia among pregnant women who have intestinal parasitic infection during current pregnancy was 227 33.33%.

**Conclusions:** The likelihood of anemia among pregnant women is approximately four times higher among pregnant women who had intestinal parasitic infection than who did not have the infection in Ethiopia

**Keywords:** Determinant of anemia, Intestinal Parasitic Infection, Meta-analysis, , Systematic reviews, Ethiopia

## INTRODUCTION

Pregnant women with Hemoglobin level less than 11 g/dl are considered to be anemic (1).

In the world, 56 million pregnant women are anemic (2) In Africa the magnitude of anemia among pregnant women was 57.1% (3)

Prevalence of anemia among pregnant women in Ethiopia is 63% and in east Africa countries 55% in Kenya, 58% in Sudan and Eritrea 55.3% (4). Different factors might leads to anemia among pregnant women. Geohelminth infections during pregnancy were associated with maternal anemia. Hookworm infection causes anemia among pregnant women and it also aggravates anemia in pregnant women (5).

Infections by helminthes leads to malnutrition, iron deficiency anemia, and increased vulnerability to other infections in infected pregnant women (6).

Other recent studies in Ethiopia have reported prevalence of anemia ranging from 16.6% in a facility-based study in Gondar, northwest Ethiopia to 56.8% in Gode town, Eastern Ethiopia [6, 7].

Prior studies in Ethiopia have reported significant associations between anemia in pregnancy and parasitic infections (e.g. schistosomiasis, hookworm infection), prior use of contraceptives, use of iron supplementation, birth spacing/ intervals, parity and gravidity, educational attainment, age, body weight, trimester of pregnancy and wealth status. [6 -18]

Despite there are many researches done on anemia among pregnant women in Ethiopia, but data on intestinal parasitic infection and anemia among pregnant women in Ethiopia is not adequate.

This systematic review and meta-analysis was conducted to assess the relationship between intestinal parasitic infection and anemia among pregnant women in Ethiopia.

## Methods

### Search approach and appraisal of studies

Studies were Searched using primary key terms of 'determinant of anemia ', 'anemia ', 'intestinal parasitic infection ', ' intestinal parasitic infection and anemia ', 'Ethiopia ' and to generate additional keywords for the search we were used the following search strategies; ' intestinal parasitic infection + anemia +pregnant women + Ethiopia through Electronic databases on reference manager software

The databases searched were PUBMED and Advanced Google Scholar

References of studies that meet eligibility criteria were used to identify similar articles

### Inclusion criteria

All Studies that were assessed the relationship between intestinal parasitic infection and anemia.

The outcome of interest was anemia

The study reported the percentage of anemia according to intestinal parasitic infection

Meet quality assessment

### Exclusion criteria

Studies that were published in languages other than English,

Included participants with anemia not dichotomized as anemia and no anemia,

Included participants with intestinal parasitic infection not dichotomized as yes and no

studies conducted not in Ethiopia were also excluded to avoid the combination of studies that were not comparable.

### Data Extraction

Three researchers were carried out the data extraction The extracted information were the name of the author, study area, , the number and percentage of anemia , the number and percentage of intestinal parasitic infection

### Risk of bias and quality assessment

To assess external and internal validity, a risk-of-bias tool was used. The tool has seven items: 1) random sequence generation (selection bias), 2) allocation concealment (selection bias), 3) blinding of participants (performance bias), 4) blinding of outcome assessment (detection bias), 5) incomplete outcome data (attrition bias), 6) selective reporting (reporting bias) and 7) other bias. All of these items are rated based on the author's subjective judgment given responses to the preceding seven items rated as low, moderate or high risk.

Three reviewers assessed independently the articles for inclusion in the review using risk-of-bias tool and guided by PRISMA checklist.

A discrepancy that would face by reviewers on selection of studies and data extraction was resolved by discussion additionally, all potential confounding variables were controlled by multivariable analysis in all included studies.

## Measures

Outcome variable pregnant women with Hemoglobin level less than 11 g/dl are considered to be anemic (1)

## Statistical analysis

The necessary information was extracted from each original study by using a format prepared in Microsoft Excel spreadsheet and transferred to Meta-essential and Revman software for further analysis.

Pooled effect size of anemia was estimated from the reported proportion of eligible studies using RevMan V.5.3 software. Forest plots were generated displaying MH odd ratio with the corresponding 95% CIs for each study. As the test statistic showed significant heterogeneity among studies Heterogeneity:  $\text{Tau}^2 = 0.13$ ;  $\text{Chi}^2 = 13.11$ ,  $\text{df} = 5$  ( $P = 0.02$ );  $I^2 = 62\%$ ) the Random effects model was used to estimate the DerSimonian and Laird's pooled effect.

## Assessment of Publication Bias

Funnel plot asymmetry and Egger's test was used to check the publication bias

## RESULT

### Selected studies

Figure 1 shows selection process of studies 120 of records identified through database searching 40 of records after duplicates removed 33 of records screened and 13 of records excluded, 20 of full-text articles assessed for eligibility and 14 of full-text articles excluded, with reasons, studies not in Ethiopia and studies not examining intestinal parasitic infection and anemia and finally 6 of studies included in quantitative synthesis (meta-analysis)




Figure 1 selection process of studies

## Characteristics of Included Studies

Six observational studies involving 2838 participants, 557 pregnant women who have anemia were included.


**Table 1 shows description of original studies included (n=6)**

S.no	Articles		Anemia	
			Yes	No
1	Lebso M, Anato A, Loha E (2017) Prevalence of anemia and associated factors among pregnant women in Southern Ethiopia: A community based cross-sectional study. PLoS ONE 12(12): e0188783. <a href="https://doi.org/10.1371/journal.pone.0188783">https://doi.org/10.1371/journal.pone.0188783</a>	Intestinal parasite infection	Yes No	53 64 108 279
2	Argaw B, Argaw-Denboba A, Taye B, Worku A, Worku A (2015) Major Risk Factors Predicting Anemia Development during Pregnancy: Unmatched-Case Control Study. J Community Med Health Educ 5: 353. doi:10.4172/2161-0711.1000353	Intestinal parasite infection	Yes No	35 77 52 397
3	Hailu et al. Determinant factors of anaemia among pregnant women attending antenatal care clinic in Northwest Ethiopia Tropical Diseases, Travel Medicine and Vaccines (2019) 5:13 <a href="https://doi.org/10.1186/s40794-019-0088-6">https://doi.org/10.1186/s40794-019-0088-6</a>	Intestinal parasite infection	Yes No	56 23 222 442
4	Fekede Weldekidan et al Determinants of Anemia among Pregnant Women Attending Antenatal Clinic in Public Health Facilities at Durame Town: Unmatched Case Control Study Anemia Volume 2018, Article ID 8938307, 8 pages <a href="https://doi.org/10.1155/2018/8938307">https://doi.org/10.1155/2018/8938307</a>	Intestinal parasite infection	Yes No	49 62 43 179
5	Getahun et al. Burden and associated factors of anemia among pregnant women attending antenatal care in southern Ethiopia: cross sectional study BMC Res Notes (2017) 10:276 DOI 10.1186/s13104-017-2605-x	Intestinal parasite infection	Yes No	21 39 7 150
6	Kebede et al. The magnitude of anemia and associated factors among pregnant women attending public institutions of Shire Town, Shire, Tigray, Northern Ethiopia, 2018 BMC Res Notes (2018) 11:595 <a href="https://doi.org/10.1186/s13104-018-3706-x">https://doi.org/10.1186/s13104-018-3706-x</a>	Intestinal parasite infection	Yes No	13 65 22 380

In all studies, the proportion of anemia among pregnant women who have intestinal parasitic infection during current pregnancy was 227 33.33%).

## Pooled effect size


The odds of anemia among pregnant women who had intestinal parasitic infection is 3.74 times higher than those pregnant women who did not have intestinal parasitic infection (ORMH = 3.74, 95% CI 2.58-5.43) Heterogeneity:  $\tau^2 = 0.13$ ;  $\chi^2 = 13.11$ ,  $df = 5$  ( $P = 0.02$ );  $I^2 = 62\%$  Test for overall effect:  $Z = 6.94$  ( $P < 0.00001$ )



**Figure 2:** Forest plot for the association between intestinal paresis infection and anemia among pregnant women in Ethiopia

### Assessment of Publication bias analysis

No publication bias was observed (Egger's test:  $p = 0.074$ , Begg's test:  $p = 0.091$ ).




Egger Regression					
	Estimate	SE	CI LL	CI UL	
Intercept	5.12	2.13	-0.36	10.61	
Slope	-0.31	0.66	-2.01	1.40	

t test	2.40
p-value	0.074

Begg & Mazumdar	
$\Delta_{x-y}$	9.00
Kendall's Tau a	0.60
z	1.69
p	0.091

Figure\_3: Funnel plot for the association between intestinal paresis infection and anemia among pregnant women in Ethiopia

The below 'Abbe plot showed that all studies effect size is above the zero effect size




### Combined effect size

Fixed effect model

Risk Ratio      2.47

Figure\_4: 'Abbe plot for the association between intestinal paresis infection and anemia among pregnant women in Ethiopia

The below Galbraith plot showed that 95% of the studies effect size lie between the two color lines and this indicates there is no outlier of effect size



Regression estimate				
	Estimate	SE	CI LL	CI UL
Intercept	0.00			
Slope	1.26	0.19	0.77	1.74

Figure\_5: Galbraith plot for the association between intestinal parestis infection and anemia among pregnant women in Ethiopia

## DISCUSSION

In this systematic review and meta-analysis, pregnant women who were infected with intestinal parasite were 3.74 times more likely to be anemic than who were not infected by any intestinal parasite and this is consistent with previous studies, (6, 19-24).

The worm in the intestine may cause intestinal necrosis and blood loss as a result of the attachment to the intestinal mucosa and chronic infections lead to iron deficiency and anemia resulting from the excessive loss of iron (25).

Therefore, an effective intervention packages need to reduce anemia among pregnant women through iron supplementation, anthelmintic treatment and dietary diversification in the study area (26)

Our finding is similar with other previous study in Ethiopia (27) this possibly happen because most anemic pregnant women who are living in Ethiopia were farmers, bare foot walking is common among Ethiopian farmers, and the chance to be exposed for soil transmitted parasite is very high. Besides this, the low environmental sanitation status may also aggravate the chance of intestinal parasite infection.

Parasitic diseases were known to play as a major contributing factor to anemia in pregnancy. For example, blood loss caused by hookworm puts mothers at high risk of iron deficiency anemia (28).

Our finding is similar with many previous studies conducted in Ethiopia and other developing countries that have shown the strong association of intestinal parasitic infection with anemia in pregnant women (23, 28-29, 30--33) Parasitic

infection has a devastating effect on the level of Hgb and causes anemia since they affect iron absorption by the intestine and consumes the red blood cells (34).

There was a strong significant association between intestinal parasitic infection and anemia in pregnant women in previous studies (35), in Southern Ethiopia (19), Ghana (34), Nigeria (36), and Venezuela (37), Durame Town Ethiopia (38), in Yirgalem and Hawassa cities, Dessie town and Canada (27, 39, 33), Shire Town, Tigray, Northern Ethiopia (40), in Shalla Woreda of Oromia region In Ethiopia (16).

This is expected as intestinal parasites, apart from their competition for nutrients, are known to cause blood loss, loss of appetite reduced motility of food through the intestine and damage to the wall of the intestine leading to mal-absorption of nutrients.

## CONCLUSION

The likelihood of anemia among pregnant women is approximately four times higher among pregnant women who had intestinal parasitic infection than who did not have the infection

### Data Availability:

All data are included in the paper.

Conflicts of Interest: The authors declare that they have no conflicts of interest.

### Authors' Contributions:

- Kaleab Tesfaye Tegegne was responsible for conceptualization, project administration, software, supervision, and development of the original drafting of the manuscript.
- Kaleab Tesfaye Tegegne, Eleni Tesfaye Tegegne, and Mekibib Kassa Tessema were participated in quality assessment of articles, methodology, validation, and screening of research papers
- All authors contributed with data analysis, critically revised the paper, and agreed to be accountable for their contribution.

### Acknowledgments

We would like to thank all the primary authors of the included articles

### Competing of interest

The authors have declared that there is no competing interest

### Funding

Not any funding received for this work

## REFERENCES

1. De Maeyer EM, Dallman P, Gurney JM, Hallberg L, Sood SK, Srikantia SG. Preventing and controlling iron deficiency anemia through primary health care, a guide for health administrators and programme managers; 1989.
2. Balarajan Y, Ramakrishnan U, Ozaltin E, Shankar AH, Subramanian SV. Anaemia in low-income and middle-income countries. *The Lancet*, 2011; 378 (9809):2123-2135
3. World health organization. Worldwide prevalence of anemia 1993-2005. WHO Global Database on Anemia. Geneva: World health organization; 2008
4. Benoist BD, McLean E, Egll I, Cogswell M (2008) Worldwide prevalence of anemia 1993-2005: WHO global database on anemia. World Health Organization, Geneva, Switzerland
5. Stephenson LS, Latham MC, Ottesen EA. Malnutrition and parasitic helminth infections. *Parasitology*. 2000;121:23-38.
6. Melku M, Addis Z, Alem M, Enawgaw B. Prevalence and Predictors of maternal anemia during pregnancy in Gondar, Northwest Ethiopia: an institutional based cross-sectional study. *Anemia*. 2014;2014
7. Alene KA, Dohe AM. Prevalence of Anemia and Associated Factors among Pregnant Women in an Urban Area of Eastern Ethiopia. *Anemia*. 2014;2014
8. Kefyalew AA, Abdulahi MD. Prevalence of anemia and associated factors among pregnant women in an urban area of eastern Ethiopia. *Anemia*. 2014;2014:561-7.
9. Desalegn S. Prevalence of anaemia in pregnancy in Jima town, southwestern Ethiopia. *Ethiop Med J*. 1993;31(4):251-8.
10. Abriha A, Yesuf ME, Wassie M. Prevalence and associated factors of anemia among pregnant women of Mekelle town: a cross sectional study. *BMC Res Notes*. 2014;7:888.



11. Bekele A, Tilahun M, Mekuria A. Prevalence of anemia and its associated factors among pregnant women attending antenatal Care in Health Institutions of Arba Minch town, Gamo Gofa Zone, Ethiopia: a cross-sectional study. *Anemia*. 2016;2016:1073192.
12. Gebre A, Mulugeta A. Prevalence of anemia and associated factors among pregnant women in north western Zone of Tigray, northern Ethiopia: a cross-sectional study. *J Nutr Metab*. 2015;2015:165430.
13. Gedefaw L, Ayele A, Asres Y, Mossie A. Anemia and associated factors among pregnant women attending antenatal Care Clinic in Wolayita Sodo Town. *Southern Ethiopia Ethiop J Health Sci*. 2015;25(2):155–62.
14. Getachew M, Yewhalaw D, Tafess K, Getachew Y, Zeynudin A. Anaemia and associated risk factors among pregnant women in Gilgel gibe dam area, Southwest Ethiopia. *Parasites Vectors*. 2012;5:296.
15. Kedir H, Berhane Y, Worku A. Khat chewing and restrictive dietary behaviors are associated with anemia among pregnant women in high prevalence rural communities in eastern Ethiopia. *PLoS One*. 2013;8(11):e78601.
16. Obse N, Mossie A, Gobena T. Magnitude of anemia and associated risk factors among pregnant women attending antenatal care in Shalla Woreda, west Arsi Zone, Oromia region, Ethiopia. *Ethiop J Health Sci*. 2013;23(2):165–73.
17. Alemu T, Umeta M. Reproductive and obstetric factors are key Predictors of maternal anemia during pregnancy in Ethiopia: evidence from Demographic and health survey (2011). *Anemia*. 2015;2015:649815.
18. Zerfu TA, Umeta M, Baye K. Dietary diversity during pregnancy is associated with reduced risk of maternal anemia, preterm delivery, and low birth weight in a prospective cohort study in rural Ethiopia. *Am J Clin Nutr*. 2016; doi:10.3945/ajcn.115.116798.
19. Lebso M, Anato A, Loha E (2017) Prevalence of anemia and associated factors among pregnant women in Southern Ethiopia: A community based cross-sectional study. *PLoS ONE* 12(12): e0188783. <https://doi.org/10.1371/journal.pone.0188783>
20. Meseret A, Bamlaku E, Aschalew G, Tigist K, Mohammed S, Yadessa O, et al. Prevalence of anemia and associated risk factors among pregnant women attending antenatal care from Gondar town, Northwest Ethiopia. *J Inter Histo*. 2013; 1(3):137±144
21. Makhoul Z, Taren D, Duncan B, Pandey P, Thomson C, Winzerling J, et al. Risk factors associated with anemia, iron deficiency and iron deficiency anemia in rural Nepal pregnant women. *Southeast Asian J Trop Med Pub Heal*. 2012; 43(3):735±746.,
22. Oueâdraogo S, Koura KG, Accrombessi KM, Bodeau-Livinec F, Mssougbojji A, Cot M, et al. Maternal anemia at first antenatal visit: prevalence and risk factors in a malaria-endemic area in Benin. *Am J Trop Med Hyg*. 2012; 87(3):418±424. <https://doi.org/10.4269/ajtmh.2012.11-0706> PMID: 22826498
23. Horque M, Horque E, Kader SB. Risk factor of anaemia in pregnancy in rural Kuwazula- Natal South Africa, implication for health education and promotion. *SA Fam Pract*. 2009; 51(1):68±72.
24. Tefera G. Determinants of anemia in pregnant women with emphasis on intestinal helminthic infection at Sher-Ethiopia Hospital, Ziway, Southern Ethiopia. *Immunology and Infectious Diseases*. 2014; 2(4):33±39
25. World health organization. Iron deficiency anaemia: assessment, prevention and control. A guide for programme managers. Geneva, 2001 (WHO/NHD/01.3)
26. Stoltzfus RJ, Dreyfuss ML, Chwaya HM, Albonico M. Hookworm control as a strategy to prevent iron deficiency. *Nutr Rev*. 1997; 55: 223±232. PMID: 9279058
27. Argaw B, Argaw-Denboba A, Taye B, Worku A, Worku A (2015) Major Risk Factors Predicting Anemia Development during Pregnancy: Unmatched-Case Control Study. *J Community Med Health Educ* 5: 353. doi:10.4172/2161-0711.1000353
28. Brooker S, Hotez PJ, Bundy DAP (2008) Hookworm-Related Anaemia among Pregnant Women: A Systematic Review. *PLoS Negl Trop Dis* 2: e291.
29. Bethony P, Brooker S, Hotez P (2006) Soil-transmitted helminth infections: ascariasis, trichuriasis, and hookworm. *Lancet* 367: 1521-1532
30. Rodríguez-Morales AJ, Barbella RA, Case C, Arria M, Ravelo M, et al. (2006) Intestinal parasitic infections among pregnant women in Venezuela. *Infect Dis Obstet Gynecol* 2006: 23125.
31. Yatch NJ, Yi J, Agbenyega T, Turpin A, Rayner JC, et al. (2009) Malaria and intestinal helminth co-infection among pregnant women in Ghana: prevalence and risk factors. *Am J Trop Med Hyg* 80: 896-901.
32. Alem M, Enawgaw B, Gelaw A, Kena T, Seid M, et al. (2013) Prevalence of anemia and associated risk factors among pregnant women attending antenatal care in Azezo Health Center Gondar town, Northwest Ethiopia. *J Interdiscipl Histopathol* 3: 137-144
33. Gyorkos TW, Gilbert NL (2014) Blood Drain: Soil-Transmitted Helminths and Anemia in Pregnant Women. *PLoS Negl Trop Dis* 8: e2912.
34. Tay KSC, Nani EA, Walana W. Parasitic infections and maternal anaemia among expectant mothers in the Dangme East District of Ghana. *BMC Res Notes*. 2017;10:

35. Hailu et al. Determinant factors of anaemia among pregnant women attending antenatal care clinic in Northwest Ethiopia Tropical Diseases, Travel Medicine and Vaccines (2019) 5:13 <https://doi.org/10.1186/s40794-019-0088-6>
36. Akinbo FO, Olowookere TA, Okaka CE, Oriakhi MO. Co-infection of malaria and intestinal parasites among pregnant women in Edo State, Nigeria. J Med Trop. 2017;19:43–8
37. Rodríguez-Morales AJ, Barbella AR, Case C, Melissa Arria M, Ravelo M, Perez H, Oscar U, Gervasio G, Rubio N, Maldonado A, Aguilera Y, Viloria A, Blanco JJ, Colina M, Hernández E, Araujo E, Cabaniel G, Benitez J, Rifakis P. Intestinal Parasitic Infections among Pregnant Women in Venezuela. Infect Dis Obstet Gynecol 2006;2006:23125
38. Fekede Weldekidan et al. Determinants of Anemia among Pregnant Women Attending Antenatal Clinic in Public Health Facilities at Durame Town: Unmatched Case Control Study Anemia Volume 2018, Article ID 8938307, 8 pages <https://doi.org/10.1155/2018/8938307>
39. S. E. Tadesse, O. Seid, Y. G. Mariam et al., “Determinants of anemia among pregnant mothers attending antenatal care in Dessie town health facilities, northern central Ethiopia, unmatched case -control study,” PLoS ONE, vol. 12, no. 3, 2017
40. Kebede et al. The magnitude of anemia and associated factors among pregnant women attending public institutions of Shire Town, Shire, Tigray, Northern Ethiopia, 2018 BMC Res Notes (2018) 11:595 <https://doi.org/10.1186/s13104-018-3706-x>