



## Utilizing Amniotic Membrane Allograft as regenerative therapy for Treating Burns

Srinivasan D<sup>1</sup>, \*Ravi Kumar Chittoria<sup>2</sup>, Kanav Gupta<sup>3</sup>

<sup>1</sup>Junior Resident (NPG), Department of Plastic Surgery, Puducherry, India – 605006

<sup>2</sup>MCh, DNB, MNAMS, FRCS (Edin), DSc, PhD (Plastic Surgery) Professor & Registrar (Academic) Head of IT Wing and Telemedicine Department of Plastic Surgery & Telemedicine JIPMER Pondicherry India – 605006

<sup>3</sup>MBBS, MS DNB General Surgery Senior Resident Department of Plastic Surgery JIPMER Puducherry India – 605006

DOI: 10.5281/zenodo.16755413

Submission Date: 24 June 2025 | Published Date: 07 Aug. 2025

\*Corresponding author: Ravi Kumar Chittoria

MCh, DNB, MNAMS, FRCS (Edin), DSc, PhD (Plastic Surgery) Professor & Registrar (Academic) Head of IT Wing and Telemedicine Department of Plastic Surgery & Telemedicine JIPMER Pondicherry India – 605006

### Abstract

Burn injuries are prevalent among all age groups and can result from thermal, scald, or electrical sources. Currently, various scaffolds are employed to enhance the healing process and minimize scar formation. Collagen serves as a scaffold, facilitating tissue regeneration and promoting the formation of new blood vessels. Additionally, scaffolds such as amniotic membrane aid in proper epithelialization and scar reduction, boasting unique anti-inflammatory and bacteriostatic properties. In our study, we utilized amniotic membrane allograft as a biological dressing for burn wounds on a infant patient's anterior chest.

**Keywords:** Amniotic membrane allograft, wound healing, regenerative therapy.

## Introduction

The process of wound healing is a biological response to various forms of injury, encompassing physical, chemical, mechanical, or thermal damage. This process typically involves several sequential phases: homeostasis, inflammation, proliferation/granulation, and remodeling/maturation<sup>1</sup>. However, deviations from the normal healing trajectory often result in stagnation during the inflammatory phase, particularly evident in cases of burns where standard healing mechanisms may be compromised<sup>2</sup>. In contemporary medicine, the utilization of scaffolds, whether natural or synthetic, has gained traction and recognition. An optimal scaffold should possess specific characteristics: appropriate physical and mechanical properties, a physiological background conducive to cellular adhesion, proliferation, and differentiation, high porosity, a favorable surface area to volume ratio, flexibility to conform to wound shapes, and ideally, biocompatibility and biodegradability<sup>3</sup>. Collagen, whether synthetic or natural, serves as a surrogate for the dermal matrix, facilitating epithelialization during wound healing<sup>4</sup>. Additionally, collagen degradation during this process contributes to neovascularization, thereby promoting angiogenesis. Amniotic membrane, a natural scaffold, exhibits distinctive properties such as anti-inflammatory, bacteriostatic, anti-fibrotic, anti-scarring, and epithelialization-promoting qualities<sup>6</sup>. Its low immunogenicity and presence of progenitor cells make it an appealing choice for scaffold applications in wound healing. Silicone functions as a barrier, reducing mechanical friction and transepidermal water loss, factors associated with infection severity. In vitro studies suggest that silicone may modulate inflammatory growth factors implicated in fibrosis and support acute wound healing, including key inflammatory markers like TNF- $\alpha$ , TGF- $\beta$ , IL-1, and IL-6.

## Materials and Methods

This research was carried out at a Tertiary Care Centre within the Department of Plastic Surgery following approval from the department's ethical committee. Informed consent was duly obtained. . The subject is a 1 year old female patient, with alleged history of accidental burns with hot milk over face, neck and chest (Fig 1). She was taken to local hospital where primary care was done and referred to JIPMER with delay of 1 hours. Fluid resuscitation was initiated as per parkland formula The patient had undergone multiple dressings of functional regenerative therapy with scaffold along with amniotic membrane and cyclical NPWT. This scaffold comprised sterile amniotic membrane, dry collagen sheet, and

silicone sheet, with the amniotic membrane layer directly contacting the wound. The amniotic membrane (fig 2) used was obtained from a freshly delivered placenta, thoroughly washed, and stored in an antibiotic solution in a refrigerator. The dressing remained in place for four days, during which time the amniotic membrane was completely absorbed, and the silicone sheet layer was also removed (fig 3).



**Fig 1. Wound at presentation**



**Fig 2. Figure showing amniotic membrane which was harvested from human placenta**





**Fig 3. Figure showing amniotic membrane which was harvested from human placenta being applied over the wound over the anterior chest region**

### **Results**

Throughout both the intraoperative and postoperative phases, the patient experienced no notable events. Upon opening the dressing on the 4th day post-operation, substantial areas of re-epithelialization and healing were observed (fig 4). Complete healing of all second-degree superficial burn wounds was achieved without any complications or adverse effects noted throughout the entire procedure after utilization of amniotic membrane regenerative therapy at the time of discharge.



**Fig 4. Figure showing partial epithilization of the wound over the anterior chest region**

### Discussion

Partial-thickness burn injuries have the potential to heal spontaneously, whereas full-thickness burns typically necessitate skin grafting to achieve complete wound closure. Historically, split-thickness skin autograft has been the preferred method for closing excised full-thickness burn wounds<sup>5</sup>. However, patients with extensive burn injuries may face challenges due to limited availability of donor sites for autograft harvesting, prompting consideration of alternative options such as skin substitutes. Engineered skin substitutes offer temporary coverage for wounds until donor sites are available for autografting or, if containing autologous cells, can provide permanent closure<sup>7</sup>. While the availability of permanent skin substitutes remains limited, advancements in tissue engineering are anticipated to yield improved models, thus enhancing the options for managing burn wounds. For instance, the commercially available Dermal Regeneration Template comprises a two-layered system. The outer layer, composed of a thin silicone film, serves as a protective

barrier against infection and regulates heat and moisture loss. The inner layer, consisting of a collagen glycosaminoglycan (GAG) thermal layer, acts as a biodegradable template facilitating dermal tissue regeneration. The porous nature of the template aids in skin regeneration by serving as a scaffold for tissue growth. Following regeneration of the dermal layer, the outer layer of the template is replaced with a thin epidermal skin graft, allowing for flexible, permanent skin regeneration and promoting faster wound healing with minimal scarring. Seeking to emulate this mechanism, we have developed an indigenous dermal regeneration scaffold using silicone sheet, dry collagen sheets, and amnion<sup>3</sup>. This cost-effective scaffold can be readily prepared and utilized, particularly in hospital settings within developing countries where the affordability of commercial regeneration templates may be a concern.

## Conclusion

The efficacy of employing a cost-effective regenerative scaffold dressing based on amniotic membrane in treating second-degree scald burns has been demonstrated in this investigation. It accelerates the overall healing process of both superficial and deep second-degree wounds to less than a week, thereby reducing hospitalization duration and infection rates. Nevertheless, further extensive multicenter, double-blinded controlled research incorporating statistical analysis is warranted.

## References

1. Haghpahanah, S., Bogie, K., Wang, X., Banks, P. G., & Ho, C. H. (2006). Reliability of electronic versus manual measurement techniques. *Archives of Physical Medicine and Rehabilitation*, 87(10), 1396-1402. <https://doi.org/10.1016/j.apmr.2006.06.011>
2. Chitoria, R. K., Kumar, P., Baljaj, S. P., Singh, A. K., & Gupta, D. K. (2014). General clinical guidelines for wound management: Redefining acronym SWCR. *Journal of SWCR*, 7(1), 2-7.
3. Shetty, R., Sreekar, H., Lamba, S., & Gupta, A. K. (2012). A novel and accurate technique of photographic wound measurement. *Indian Journal of Plastic Surgery*, 45(2), 425-429. <https://doi.org/10.4103/0970-0358.101095>
4. Majeske, C. (1992). Reliability of wound surface area measurement. *Physical Therapy*, 72, 138-141.
5. Mayrovitz, H. N., & Soontupe, L. B. (2009). Wound area by computerized planimetry of digital images. *Advances in Skin & Wound Care*, 22(5), 222-229. <https://doi.org/10.1097/01.ASW.0000354924.16816.6b>
6. Pandey, S., Chitoria, R. K., Mohapatra, D. P., et al. (2016). Application of digital planimetry: A novel technique of wound measurement in diabetic foot ulcer. *Dermatology International*, 21(6), 343-331. <https://doi.org/10.1159/000452073>
7. Pires, I. M., & Garcia, N. M. (2015). Wound area assessment using mobile application. In *Proceedings of the International Conference on Biomedical Electronics and Devices - Volume 1: SmartMedDev* (pp. 271-282). Lisbon, Portugal: IEEE. <https://doi.org/10.1109/BIOMED.2015.7059392>

## CITATION

Srinivasan D, Chittoria, R. K., & Kanav G. (2025). Utilizing Amniotic Membrane Allograft as regenerative therapy for Treating Burns. In *Global Journal of Research in Medical Sciences* (Vol. 5, Number 4, pp. 74–78). <https://doi.org/10.5281/zenodo.16755413>